
 Europäisches Patentamt
 European Patent Office
 Office européen des brevets

(11) Publication number:

0 189 972
A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 86300089.9

(61) Int. Cl.⁴: C 07 D 279/06

(22) Date of filing: 08.01.86

(30) Priority: 14.01.85 GB 8500863

(43) Date of publication of application:
06.08.86 Bulletin 86/32

(64) Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

(71) Applicant: FINE ORGANICS LIMITED
Teesside Site Seal Sands
Middlesbrough Cleveland, TS2 1UB(GB)

(72) Inventor: Hollowood, John
Beech Croft
Newton York YO6 5TU(GB)


(72) Inventor: Jackson, Arthur
Glebe House
Washington Tyne & Wear(GB)

(72) Inventor: Heyes, Graham
25 Sutton Street
Durham(GB)

(74) Representative: Harrison, Michael Robert et al,
Urquhart-Dykes & Lord 5th Floor Tower House Merrion
Way
Leeds, LS2 8PA(GB)

(64) Preparation of thiazine derivatives.

(67) A method for the preparation of a thiazine derivative,
having insecticidal properties, of the formula



where each R is hydrogen or an appropriate aromatic or aliphatic substituent and X is hydrogen, halogen or lower alkyl, the method comprising reacting together a sulphur donor, a compound of the formula $Y_2C=CXNO_2$, where Y is halogen (preferably chlorine) or another appropriate leaving group and a compound of the formula



EP 0 189 972 A1

0189972

Title



Preparation of thiazine derivatives.

Introduction

This invention relates to methods of preparing thiazine derivatives which have insecticidal activity.


5. Tetrahydro-2-(nitromethylene)-2H-1,3-thiazine (I) possesses broad spectrum insecticidal activity, being particularly active against lepidopterous larvae on plants. It is also useful as an intermediate in the synthesis of more stable but equally active insecticides e.g., the oxime of 5,6-dihydro-4H-1,3-thiazine-2-carboxaldehyde (II).

10




The two known synthetic methods for preparing compound I are outlined below:-

Route A



$$\begin{array}{c}
 \text{HN} \quad \text{S} \\
 | \quad // \\
 \text{C} \\
 \backslash / \\
 \text{CH}_2\text{NO}_2 \\
 | \\
 \text{CO}_2\text{Et}
 \end{array}
 + (\text{CH}_3)_2\text{SO}_4 \longrightarrow
 \begin{array}{c}
 \text{N} \quad \text{S} \\
 | \quad // \\
 \text{C} \\
 \backslash / \\
 \text{SCH}_3
 \end{array}
 + \text{ZnCl}_2 + \text{CH}_2\text{NO}_2
 \begin{array}{c}
 | \\
 \text{CO}_2\text{Et}
 \end{array}$$


$$\xrightarrow{\hspace{1cm}}
 \begin{array}{c}
 \text{HN} \quad \text{S} \\
 | \quad // \\
 \text{C} \\
 / \quad \backslash \\
 \text{O}_2\text{N} \quad \text{CO}_2\text{Et}
 \end{array}
 \xrightarrow[\text{(ii) HOAc}]{\text{(i) NaOH}}
 \begin{array}{c}
 \text{HN} \quad \text{S} \\
 | \quad // \\
 \text{C} \\
 \backslash / \\
 \text{CHNO}_2
 \end{array}$$
$$\text{Cl}^{\ominus} \text{NH}_3^{\oplus} (\text{CH}_2)_3 \text{SH} + \text{Cl}_2 \text{C} = \text{CHNO}_2$$


Route A suffers from the disadvantage that the starting material 3-aminopropane thiol hydrochloride is expensive and the yield by this route is only moderate.

Although the starting material for Route B, tetrahydro-1,3-thiazine-2-thione, is relatively inexpensive the ethyl nitroacetate required in the second stage of the synthesis is not available in commercial quantities and this rules out this procedure for manufacturing on an industrial scale.

[illegible]
$$\text{H}_2\text{NCR}_2\text{CR}_2\text{CR}_2\text{OSO}_3\text{H}$$

The sulphur donor may be any suitable source of sulphur, for example, sulphur itself, a sulphide, a hydrosulphide or hydrogen sulphide. Preferably, the sulphur donor is an ammonium or alkali metal sulphide or hydrosulphide, for instance, the alkali metal sulphide sodium sulphide.

$$\text{H}_2\text{NCH}_2\text{CH}_2\text{CH}_2\text{OSO}_3\text{H} + \text{Na}_2\text{S} + \text{Cl}_2\text{C}=\text{CHNO}_2$$


The starting aminopropylsulphate is easily prepared from 3-amino-propanol and sulphuric acid. The preferred reaction referred to above proceeds at room temperature and this would appear to rule out a reaction mechanism involving the formation of the aminopropanethiol since it is known that the formation of the thiol only occurs at an appreciable rate at temperatures greater than 80°C. It may be that the reaction mechanism involves an initial attack by the sulphide anion on the nitroethene in a Michael type addition. The resultant thiol anion may then displace the sulphate ion in the aminopropyl sulphate. Ring closure with the elimination of two molecules of hydrogen chloride then occurs to give the product.

Because of the insolubility of the dihalonitroethylene in water it is preferred to use a cosolvent. Preferably, the cosolvent is a water immiscible organic solvent such as benzene, toluene or ethylene dichloride.

Preferably, the reaction is conducted at a temperature in the range from 0-80°C although products formed at the higher temperatures are relatively more contaminated and more difficult to purify.

EXAMPLE

An example of a method in accordance with the present invention will now be described, by way of example only.

To a stirred flask fitted with a condensor were charged water (200 ml), toluene (200 ml), aminopropyl sulphate (100 gms), sodium sulphide (78 gms/60% active) and 1,1-dichloro-2-nitro ethylene (81.1 gms). The mixture was heated to 60°C and the pH maintained between 6-8 by gradual addition of sodium hydroxide solution to absorb the hydrochloric acid liberated.

When the reaction was complete (about 1 hour) the

0189972

- 5 -

reaction solution was cooled, the aqueous layer separated off and the pH adjusted to 5.5 with acetic acid.


The aqueous layer was then extracted with methylene chloride (3 times 100 ml). The methylene chloride was distilled out and the product recrystallised from isopropanol to give tetrahydro-2-(nitro methylene)-2H-1,3-thiazine having a melting point of 73-76°C.

CLAIMS:


1. A method for the preparation of a thiazine derivative of the formula

5

10



where each R is hydrogen or an appropriate aromatic or aliphatic substituent and X is hydrogen, halogen or lower alkyl, characterised in that the method comprises reacting together a sulphur donor, a compound of the formula $Y_2C=CXNO_2$ where Y is halogen (preferably chlorine) or another appropriate leaving group and a compound of the




2. A method according to claim 1 characterised in that each R is independently hydrogen or lower alkyl.
3. A method according to claim 1 or claim 2 characterised in that the sulphur donor is an ammonium or alkali metal sulphide or hydrosulphide.
4. A method according to any of the preceding claims characterised in that the sulphur donor is sodium sulphide.
5. A method according to any of the preceding claims characterised in that Y is chlorine and X is hydrogen.
6. A method according to any of the preceding claims characterised in that the reaction is conducted in the presence

of water and a cosolvent immiscible with water.


7. A method according to claim 6 characterised in that the cosolvent is benzene, toluene or ethylene dichloride.

8. A method for the preparation of a compound of the formula



characterised in that the method comprises reacting together sodium sulphide, a compound of the formula

15 $\text{Cl}_2\text{C}=\text{CHNO}_2$ and a compound of the formula





European Patent
Office

EUROPEAN SEARCH REPORT

0189972

EP 86 30 0089

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Y	EP-A-O 127 413 (FINE ORGANICS LTD.) * Claims *	1-8	C 07 D 279/06
Y	EP-A-O 115 323 (UBE INDUSTRIES LTD.) * Claims *	1,2,5-8	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			C 07 D 279/00 C 07 D 277/00 C 07 D 281/00
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 11-04-1986	Examiner CHOULY J.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.